ACM-Aminograin

agro consulting del mediterráneo s.l.

Denomination type: AMINO ACIDS MICRONUTRIENT. Group 4.1.01
ORDER AAA / 2564/2015, 27 November, amending Annexes I, II, III, IV and VI - R. D. 506/2013, June 28, about fertilizer products.

AMINOGRAM

Aspartic acid, Threonine, Arginine, Alanine, Cysteine, Tyrosine Glutamic acid: 50%, Serine, Histidine, Glycine, Valine, Methionine, Phenylalanine, Isoleucine, Leucine, Lysine, Proline.

FEATURES

ACM-Aminograin helps to overcome stressful periods. The product improves vegetative and fruiting growth. Production and quality in both fibre and grain are increased. It reduces the negative effect of herbicide treatments. It prevents the decrease in vegetative growth caused by the herbicide. It encourages vegetative growth and plant development.

DOSE AND APPLICATIONS



COTTON

Dose:
1–1.25 L/Ha.

21 days after plastic cover is removed. Beginning of flowering. After capsules shape formation



RICE AND COCOA

Dose:
1–1.50 L/Ha.

Rice. At the control stage .At pre-flowering stage. Do apply the treatments with herbicides or fungicides.

Cocoa. At the maximum vegetative growth moment.



WHEAT, RYE AND OATS

Dose:
1–1.25 L/Ha.

Apply at the flowering starts in combination with herbicides and fungicides.



CORN,SUNFLOWER AND SORGHUM

Dose:
1–1.55 L/Ha.

Apply when 2 to 4 leaves appear. Do repeat when 6 to 8 leaves appear. At the starting of the bloom.

GUARANTEED RICHNESS
in % w/w

Free Amino acids from vegetal origin: 8%
Total Nitrogen (N): 5%
Organic Nitrogen: 2,5%
Ammonia Nitrogen: 2,5%
Boron (B): 0,1%
Iron (Fe): 0,1%
Manganese (Mn): 0,45%

Copper (Cu): 0,1%
Zinc (Zn): 0,30%
Molybdenum (Mo): 0,02%
pH: 5-6
Density: 1,20 g/cm^3
Color: Dark brown

RESIDUE ZERO

CAUTIONS

The recommendations and information that we provide are the result of wide and severe test. However, in its use, many factors affecting out of our control may appear such as weather, preparing broths , applications, etc.). The company guarantees the composition, formulation and content. User will be responsible for damage caused by fault of efficacy, toxicity, waste, etc. Not to follow total or partial label directions. Store in a dry and protected from extremes temperatures (5 ºC-35 ºC) place. Keep out of reach of children. Keep away from foodstuffs, beverages and feed. In case of eyes or skin contact, wash immediately with plenty of water.

BIOESTIMULANTS | ELEMENT CORRECTORS | SOIL AMENDMENTS | BASIC SUBSTANCES | OTHERS PRODUCTS