ACM-Potasio LQ

agro consulting del mediterráneo s.l.

Designation Type: NK fertilizer solution 3-30. C.2.9. Regulation (EC) No 2003/2003. EC fertilizer

FEATURES

ACM-Potasio LQ is a rich potassium nutrient with anti-blocking agent. Special formula that enables its absorption and efficacy. Avoid locking and interactions with other nutrients. It enables the formation of sugars and oils. It regulates the transport of water in the plant.

DOSE AND APPLICATIONS



FOLIAR
Foliar spray.
Dose:
300 - 500 cc/Hl, perform 3 to 5 applications during the fattening period and fruit development.



FRUiT AND CITRUS TREES
Drip irrigation.
Dose:
8 – 12 L/Ha, perform 2 to 3 applications during the development phase and fattening fruit.



VINEYARD AND OLIVE TREE
Drip irrigation.
Dose:
8 -12 L/Ha, perform 2 to 3 applications during the development phase and fattening fruit.



HORTICULTURAL CROPS
Drip irrigation.
Dose:
6-10 L/Ha. Applications will start 1-2 weeks before the first cut repeating every 10 days.



POTATOES AND BEETS
Drip irrigation.
Dose:
8-12 L/Ha. Make 2 applications during the fattening period.



FLOWERS AND ORNAMENTALS
Drip irrigation.
Dose:
1-1,5 cc/m^2. Apply 2 weeks before cutting or growing plant.



CONTINUOUS FERTILIZATION
Drip irrigation.
Dose:
0,75 – 2 L /m^3 in irrigation water.



HYDROPONICS
Drip irrigation.
Dose:
30 - 60 cc/m^3 in irrigation water.

GUARANTEED RICHNESS in % w/w

Potassium (K_2O) soluble in water: 30,0%
Total Nitrogen (N): 3,0%
Ureic Nitrogen (N): 3,0 %
Chelating agent: EDTA 1,0%
Appearance: Liquid Soluble (LS)
Density: 1,45–1,55 g/cm^3
pH (water solution 1%): 12-13

RESIDUE ZERO

CAUTIONS
The recommendations and information that we provide are the result of a wide and severe test. However in its use many factors out of our control may appear such as weather, preparation of broths, applications, etc. The company guarantees the composition, formulation and content. The product is compatible with the most commonly fertilizers and pesticides used in agriculture. However due to diversity of other products, it is recommended to perform a previous compatibility test and select the products to be applied. User shall be liable for damage caused by lack of efficacy, toxicity, waste, etc., not to follow label instructions. Store in a dry and protected from extreme temperatures (5 °C-35 °C) place. Keep out of reach of children. Stay away from food, drink and feed. In case of eyes and skin contact wash immediately with plenty of water.

BIOESTIMULANTS | ELEMENT CORRECTORS | SOIL AMENDMENTS | BASIC SUBSTANCES | OTHERS PRODUCTS